



HIGH SIERRA HIKERS ASSOCIATION

PO Box 8920 SO. LAKE TAHOE CA 96158

September 6, 2006

Superintendent
Yosemite National Park
Attn: Tuolumne Planning
P.O. Box 577
Yosemite, CA 95389

SUBJECT: SCOPING COMMENTS ON TUOLUMNE RIVER PLAN AND TUOLUMNE MEADOWS PLAN

Dear Superintendent,

The High Sierra Hikers Association (HSHA) is a nonprofit public-benefit organization that seeks to inform and educate its members, public agencies, and the general public about issues affecting hikers and the High Sierra. Many of the HSHA's members visit the Tuolumne Meadows and Tuolumne River areas for hiking, camping, backpacking, horse packing, and other recreational pursuits. Following are our scoping comments on the Tuolumne River Plan and the Tuolumne Meadows Plan. Please place a copy of this letter in the project record for both plans.

General Comments

The HSHA is very concerned about the ongoing (and increasing) adverse impacts in these areas due to commercial stock animal usage and the High Sierra Camps. These planning processes should be used to end—once and for all—the impairment of park, wilderness, and wild & scenic river resources and values resulting from these high-impact activities. Following are our specific comments:

High Sierra Camps (HSCs)

The HSHA is especially concerned with the Glen Aulin High Sierra Camp (HSC), and the non-wilderness HSC at Tuolumne Meadows in conjunction with the Tuolumne Meadows Lodge. These aged and ugly facilities have a significant negative impact on the Tuolumne River corridor and on the environment of Tuolumne Meadows. For example, all the by-products of human occupancy are produced at the Glen Aulin camp: sewage (human body wastes), “gray water”

from showers, grease and detergent from kitchens. But there are no wastewater or sewage treatment plants. Wastewater and sewage from these developments pollutes the meadows, soils, and waters of Yosemite National Park.

Congress specifically recognized this threat to Yosemite when it passed the California Wilderness Act of 1984. That Act, signed by President Reagan, bestowed formal wilderness designation upon much of the Yosemite backcountry. The Congress allowed the HSCs to temporarily remain, but stated:

“...If and when it occurs that the continued operation of these facilities...results in an increased adverse impact on the adjacent wilderness environment (including increased adverse impact on the natural environment within the enclaves themselves), the operation of these facilities shall be promptly terminated, the facilities removed, the sites naturalized, and in the procedure set forth by section 9 of the bill, the areas promptly designated as wilderness.”

The HSCs are an anachronism—an out-of-date holdover from the bad old days of the 1920s through the early 1960s, when more development and more commercialism were considered desirable and beneficial. One way to look at the HSCs today is this: If the NPS were to propose establishing an HSC in the Yosemite backcountry at the present time, the project would never get off the ground. It would violate the Wilderness Act, it would violate the California Wilderness Act, and it wouldn't have a ghost of a chance of surviving an honest NEPA process. That being so, why should not the existing HSCs be abolished? Fifty years ago, no one talked about environmentalism. Now we have a federal agency, the EPA, and all and sundry declare themselves to be in favor of environmental protection. It is thus time for the National Park Service to catch up with the times (and pay attention to its Organic Act) by choosing preservation of park resources, scenery, wilderness character, and wild river values over ongoing exploitation and impairment.

The Vogelsang HSC does not itself lie within the Tuolumne River watershed or Tuolumne Meadows area, but its very existence nearby has a substantial adverse impact on both river and meadows. The trail from Tuolumne Meadows to Vogelsang HSC, like all trails traversed by the HSC supply trains, is battered and polluted, featuring flies and stench and dust. One is not out of sight of manure for the entire seven miles. Significantly, a recent study by scientists from the University of California (U.C. Davis Medical School) has documented that the Tuolumne River is being polluted in this area, and concluded that: “*pack animals are most likely the source of coliform [bacteria] pollution*” (Derlet and Carlson 2006, copy enclosed).

The same is true of the trail to Glen Aulin, a camp that can support a maximum of 32 people. For the sake of those 32, dozens of people every day—and during the course of an entire season, thousands—are inconvenienced and offended by the disgusting condition of the trail and the pollution of surrounding park lands.

The House Committee Report prepared for the 1984 California Wilderness Act also stated:

“...Because of the importance of continuing monitoring and assessment of this situation, immediately upon enactment of this bill into law, the Secretary of the Interior should document current baseline operational and environmental impact conditions of all of these facilities [HSC camps], and he should also, within one year of the date of enactment, report in writing to the relevant committee of the House and Senate, his findings and recommendations as to this matter. Annual assessments of this situation should thereafter be made by the Secretary to assure continued monitoring of conditions.” (House Committee Report No. 98-40)

Has the Park Service at Yosemite prepared the baseline reports and conducted the annual monitoring reports as requested by Congress? If such reports do exist, they should be made public at once, and the findings fully evaluated in these planning processes.

The HSCs at Glen Aulin and Vogelsang are classified as “potential wilderness additions,” which, by law, must be treated and managed essentially the same as wilderness. (*See* the California Wilderness Act of 1984, Section 9.) However, despite the ongoing and increased impacts of the HSCs, and the clear direction from Congress, we are aware that the NPS has made ongoing efforts to hide the impacts of these facilities from Congress and the public, and has illegally continued to use nonconforming methods (i.e., helicopters) to maintain the HSCs and to construct new developments (i.e., sewage mounds, toilets, etc.) at the HSCs. Congress specifically directed that:

“Helicopter use for routine nonemergency purposes associated with visitor use is a questionable activity in national park system wilderness areas and should be eliminated within designated national park system wilderness.” (House Committee Report No. 98-40, at p. 51)

In sum, all three of the HSCs discussed above (including the Tuolumne Meadows Lodge) should be subject to site-specific Environmental Impact Statements (EISs) as part of the Tuolumne River/Meadows planning process(es). This has not been done before, and is necessary to illuminate the scope and nature of the substantial environmental impacts of those facilities. Significant issues include, but are not limited to: (1) impaired scenery; (2) degraded trails; (3) pollution of surface and ground waters by sewage and wastewater produced at the HSCs; (4) pollution of surface waters by manure (bacteria, etc.) produced by pack animals that service the camps; (5) harm to wildlife that come in contact with sewage, kitchen/bath wastes, and human food sources; (6) harm to native songbirds due to proliferation of brown-headed cowbirds; etc.

Given the above, your two planning processes for the Tuolumne Meadows and River should include and adopt alternatives that will permanently remove all three of the HSCs discussed above, restore the sites, and propose that the potential wilderness additions at Glen Aulin and Vogelsang be designated as wilderness as intended by Congress in the California Wilderness Act (see that Act, Section 9; and House Committee Report No. 98-40).

Commercial Packstock Enterprises

The use of stock animals can be legitimate, appropriate, and even necessary for certain recreational and/or administrative purposes. We want to make clear at the beginning that we do **not** advocate or suggest the complete elimination of recreational or administrative stock use from Tuolumne Meadows or the Tuolumne River corridor. Our primary concern is that **the NPS must acknowledge and substantially reduce the many adverse impacts that are occurring due to the currently excessive and poorly controlled activities of commercial stock enterprises.**

We are aware that commercial packstock activities and impacts have increased **substantially** in recent years in both the Tuolumne River corridor and the Tuolumne Meadows area. Your planning process should begin by producing a complete disclosure of the increases in stock use, facilities, and impacts that have occurred over the past few decades. Then, your plans should significantly reduce/control commercial stock use to avoid the identified impacts, and incorporate definitive limits to prevent future harmful increases in commercial stock enterprises.

Quotas and Permits for Commercial Stock Outfitters

The Yosemite backcountry, including portions of the Tuolumne River Wild & Scenic River corridor, is so popular that quotas on its use have been implemented to prevent unacceptable impacts. We support the implementation of restrictions designed to protect park, wilderness, and wild & scenic river values. However, we remain concerned that commercial outfitters are allowed easy access when the general public is turned away due to use quotas.

A fundamental tenet of environmental science that must be acknowledged is that one horse (or mule) can produce ***at least*** as much impact as several people (see references below). Your management plans for the Tuolumne River/Meadows should state clearly that: (1) Commercial stock use of Yosemite National Park is a privilege—not a right, and (2) Commercial stock use shall not be given priority over private foot travel. **Wherever rationing (i.e., a quota system) is necessary, commercial stock use shall be reduced to maximize the number of people allowed to enjoy the area.**

In addition, all commercial outfitters (or their clients) should have to wait in line with the rest of the public to obtain wilderness reservations and permits. Commercial packstock enterprises should **never** be allowed to issue their own permits to conduct commercial operations in Yosemite National Park. (This is a ridiculous notion, and one that illustrates the unfair special treatment that commercial packers receive from land managers in some areas.)

Impacts of Recreational Stock Use

Parties traveling with stock animals have **much** greater impact on park, wilderness, and wild & scenic river resources and values than groups traveling on foot. The disproportionate amount of impact created by stock users must be much more limited and much better controlled.

Impacts to meadows, stream zones, wetlands, and lakeshores. Numerous studies have documented adverse impacts to meadows caused by stock animals used for recreation (Cole 1977, Merkle 1963, Nagy and Scotter 1974, Neuman 1990 & 1991a-b, Strand 1972, Strand 1979a-c, Sumner and Leonard 1947, Weaver and Dale 1978).

Trampling and grazing by livestock are known to increase soil compaction and to contribute to streambank erosion, sedimentation, widening and shallowing of channels, elevated stream temperatures, and physical destruction of vegetation (Behnke and Raliegh 1978, Bohn and Buckhouse 1985, Kauffman and Krueger 1984, Kauffman et al. 1983, Siekert et al. 1985). Streambanks and lakeshores are particularly susceptible to trampling because of their high moisture content (Marlow and Pogacnik 1985). Unstable streambanks lead to accelerated erosion and elevated instream sediment loads (Duff 1979, Winegar 1977).

In sum, the impacts of recreational stock animals on meadows, streams, wetlands, and lakeshores are substantial, and need to be addressed in this planning process.

Impacts due to invasive weeds. The role of herbivores in dispersing weeds is now well established. Seeds can be spread from one location to another by attachment to the bodies of animals (epizoochory) or by being ingested and later excreted (endozoochory). (See, for example, Fenner 1985, Hammit and Cole 1987, Harmon and Kiem 1934, Heady 1954, Janzen 1982, Ridley 1930.) Many native herbivores have been shown to be effective seed dispersers. In addition, domestic stock animals such as cattle, sheep, pigs, and horses have all been shown to pass viable seeds through their intestinal tracts. (See, for example, Harmon and Kiem 1934, Harper 1977, Heady 1954, Janzen 1981 and 1982, McCully 1951, Piggin 1978, St John-Sweeting and Morris 1991, Welch 1985.) A detailed review of the scientific literature regarding the spread of weeds by domestic livestock (cattle, sheep, and horses) concluded:

“Recent research showing that livestock significantly increase invasions by nonindigenous plants in the western U.S. is persuasive. Similar results were found in all western states and for nearly every introduced species that has been studied. Although many of these studies would have benefited from both better replication and more recent research techniques, the pattern of evidence is overwhelming.” (Belsky and Gelbard 2000)

Numerous other reports document specifically that recreation livestock (i.e., horses, mules, etc.) can and do spread exotic weeds. (See Benninger 1989, Benninger-Truax et al. 1992, Campbell and Gibson 2001, Hammit and Cole 1987, Harmon and Kiem 1934, Janzen 1981 and 1982, Landsberg et al. 2001, Quinn et al. 2006, Weaver and Adams 1996.) For example, several reports show that horses can excrete viable seeds for many days or even weeks after ingestion. (See, for example, Janzen 1981, and St John-Sweeting and Morris 1991.) Hammit and Cole (1987) state that horse manure is a major source for exotic seeds in wilderness recreation areas. Campbell and Gibson (2001) found that “seeds transported via horse dung can become established on trail systems,” and that weed seeds found in horse manure had become established along trails used by horses, but not along trails that weren’t used by horses. Weaver and Adams (1996) documented “substantial overlap in the weed species germinated from horse manure and the

weeds present along trails used by horses.” After reviewing all available scientific evidence, Landsberg et al. (2001) concluded that “concerns about dispersal of weeds by horses are legitimate.”

Invasive (i.e., weed) species have been specifically identified—at the national level—as one of the four greatest threats to our national forests.¹ The spread of invasive weeds has also been identified by the Regional Forester as an urgent issue that needs to be addressed in all Forest Service activities in California.² Current direction requires Forest Service units neighboring Yosemite to address these issues. For example, specific Standards and Guidelines applicable to neighboring Forest Service lands include³:

42. Encourage use of certified weed free hay and straw. Cooperate with other agencies and the public in developing a certification program for weed free hay and straw. Phase in the program as certified weed free hay and straw becomes available. **This standard and guideline applies to pack and saddle stock used by the public, livestock permittees, outfitter guide permittees, and local, State, and Federal agencies.**
43. Include weed prevention measures, as necessary, when amending or re-issuing permits (**including, but not limited to, livestock grazing, special uses, and pack stock operator permits**).

As outlined above, scientists have (in the past five to seven years) documented “overwhelming” evidence that domestic livestock (including horses, mules, etc.) can and do spread harmful weeds. This relatively new issue has never been adequately evaluated by the NPS at Yosemite. Therefore, your plans for the Tuolumne River and Tuolumne Meadows areas should address the issues of weeds and plant pathogens that may be spread by domestic stock animals.

This would include, at minimum, a range of reasonable alternatives for mitigating the potential for spread of weeds and plant pathogens, such as: (1) prohibiting all grazing by domestic stock (to minimize the free-roaming of stock animals and dispersion of seeds across the landscape via epizoochory and endozoochory); (2) requiring stock users to feed their animals weed-free forage for at least several days before entering the park (in order for stock animals to excrete viable weed seeds before entering Yosemite); and (3) cleaning stock coats and hooves before entering the park (to minimize the potential for epizoochory).

Given the above-described impacts, your management plans for the Tuolumne River/Meadows should include the following elements to mitigate these impacts:

- No grazing by recreation livestock should be permitted. Stock users should be required to carry feed for their animals, as is required in many other national parks. Certified weed-free feed should be required to minimize the spread of weeds. This is consistent with the

¹ See <http://www.fs.fed.us/projects/four-threats/>

² See <http://www.fs.fed.us/r5/noxiousweeds/>

³ See <http://www.fs.fed.us/r5/snfpfa/final-seis/rod/appendix-a/standards-guidelines/forest-wide.html>

biocentric approach described in Hendee and others (1990).

- Lower group size limits for stock parties should be adopted to mitigate the greater impact of stock on park resources and wild & scenic river values (see below for detailed discussion of group size limits).

Trail damage by stock animals. When compared to hikers, stock parties cause substantially greater impacts to trails (Dale and Weaver 1974, Frissell 1973, Kuss et al. 1986, Laing 1961, McQuaid-Cook 1978, Trottier and Scotter 1975, Weaver and Dale 1978, Weaver et al. 1979, Whitson 1974, Whittaker 1978, Wilson and Seney 1994).

Whitson (1974) provides a good discussion of how horse impact differs from hiker impact. Dale and Weaver (1974) observed that trails used by horses were deeper than trails used by hikers only. Trottier and Scotter (1975) documented deterioration of trails used by large horse parties. Weaver and Dale (1978) found that horses caused significantly greater trail damage than hikers. Whittaker (1978) concluded that horses significantly increased the potential for severe erosion by churning soil into dust or mud. Weaver et al. (1979) found that horses caused more trail wear than both hikers and motorcycles. After reviewing the available literature, Kuss et al. (1986) concluded that: “*Pack stock and horse travel is considerably more damaging to trails than hiking.*” Recent research (Wilson and Seney 1994) has confirmed these earlier studies, concluding that “*horses produced significantly larger quantities of sediment compared to hikers, off-road bicycles, and motorcycles.*”

To mitigate these impacts of stock use, your Tuolumne River/Meadows management plan(s) should include the following elements:

- Groups using stock should be limited to ten or fewer animals per party (as suggested by Cole 1989 & 1990).
- To allow reasonable access for stock users, and to reduce the impacts of stock use on trails, some trails should be designated and maintained to withstand stock travel. Proper maintenance of these trails (and reconstruction where necessary) may reduce (but not offset) the impacts of stock travel.
- A network of “foot travel only” trails must be designated so that hikers can enjoy a stock-free experience. These trails should be maintained for *foot travel only*. Funds saved by designating a network of “foot travel only” trails could be used for intensive maintenance of the stock trails (see Cole [1990], p. 461).

Water quality impacts of stock animals. Stock urine and manure contribute to eutrophication of streams and lakes (Stanley et al. 1979). Such impacts are a significant concern in the oligotrophic aquatic environments of Yosemite National Park. Livestock manure can also pollute water with harmful bacteria and other organisms such as *Giardia* and *Cryptosporidium*, which are pathogenic to humans and other animals. (See, for example, Derlet and Carlson 2002 and 2006).

Some stock users continue to claim that the strains of *Giardia* and *Campylobacter* spread by domestic livestock are not infective to humans. This is wishful thinking. For example, their argument that humans cannot contract *Giardia* from stock animals hinges on a single inconclusive study conducted on domestic cats. The cross-transmission of enteric pathogens from stock animals is certainly not fully understood. However, there is an increasing body of evidence showing that pathogenic bacteria, protozoa such as *Giardia* and *Cryptosporidium*, and other harmful pathogens can be spread from stock animals to humans (Bemrick 1968, Blaser et al. 1984, Buret et al. 1990, Capon et al. 1989, Davies and Hibler 1979, Derlet and Carlson 2002, Derlet and Carlson 2006, Faubert 1988, Isaac-Renton 1993, Kasprzak and Pawlowski 1989, Kirkpatrick and Skand 1985, Kirkpatrick 1989, LeChevallier et al. 1991, Manahan 1970, Manser and Dalziel 1985, Meyer 1988, Rosquist 1984, Saeed et al. 1993, Strandén et al. 1990, Suk 1983, Suk et al. 1986, Taylor et al. 1983, Upcroft and Upcroft 1994, Weniger et al. 1983, Xiao et al. 1993).

Specifically, Derlet and Carlson (2002) found pathogenic organisms in 15 of 81 manure samples collected from pack animals along the John Muir Trail. This documents that about twenty percent of the manure piles in the Sierra contain potentially pathogenic organisms (i.e., organisms that may cause disease in humans). Pack animal manure collected in the Tuolumne River/Meadows areas contained pathogenic bacteria as well as *Giardia*. Derlet and Carlson (2006) also found pathogenic bacteria in surface waters of the Tuolumne River watershed, and concluded that “***pack animals are most likely the source of coliform [bacteria] pollution.***”

Your environmental document must evaluate and disclose the effects of domestic animal wastes on the environment, and your management plan(s) should include the following elements to minimize the amount of animal waste that reaches water courses:

- Campsites for stock users should be designated away from water, on level and dry sites. Stock users should be required to camp at these designated sites, and to keep their animals tied at all times when not in use. This will require stock users to carry feed for their animals, as is required in many other national parks. Managers should carefully select and designate campsites and hitching sites for such use (see Cole [1990], pp. 457-462).
- Stock users should be required to use other management tools (i.e., use of portable electric fencing when watering stock, diapers on horses, etc.) to prevent pollution of surface waters by livestock manure. (See enclosed report “Horses in Diapers Help Mexico’s Beach Cleanup.” This report documents the feasibility of requiring diapers on horses to prevent the spread of diseases found in horse manure. Horse diapers are commercially available and have been accepted around the world.⁴

⁴ See <http://www.equisan.com.au/>

In addition, your environmental document must acknowledge not only the State's specific water quality standards, but also the state/federal anti-degradation requirements.⁵ Significantly, the waters of Yosemite National Park are high quality waters that are eligible for designation as Outstanding National Resource Waters. The federal and State anti-degradation requirements clearly apply. Specifically, the National Park Service must comply with the California State Water Board's Resolution No. 68-16, which requires that existing high quality waters be fully protected, unless very specific formal findings are made. In this case, neither the Central Valley Regional Water Quality Control Board, the California State Water Resources Control Board, nor the U.S. Environmental Protection Agency has ever made the overriding findings necessary to allow degradation of water quality from the High Sierra Camps or the commercial stock enterprises that operate within Yosemite. Therefore, because the degradation and pollution of water resulting from both the High Sierra Camps and the commercial pack & saddle stock enterprises are *controllable*, that degradation and pollution must be fully prevented (unless the findings required by Res. 68-16 are formally made).

Impacts of brown-headed cowbirds. The operation of livestock pack stations, stables, and corrals (i.e., stock holding areas) is contributing to the demise of songbird populations in the Sierra Nevada by creating artificial habitat for the parasitic brown-headed cowbird. Cowbirds are obligate brood parasites that can significantly impact native passerine species. One study in the northern Sierra found that up to 78 percent of warbler nests are parasitized by cowbirds, resulting in significant decreases in the reproductive success of those species (Airola 1986). Elsewhere in the Sierra, individual female cowbirds have been reported to lay an average of 30 eggs per season (Fleischer et al. 1987). These high rates of parasitism and fecundity by cowbirds indicate that significant local impacts occur wherever cowbird populations are present. Habitat modifications, pack stations, corrals, and the presence of livestock throughout the Sierra may contribute significantly to regional declines in songbird populations (Graber 1996). A detailed literature review on cowbird impacts is enclosed and incorporated by reference. The impacts of stock holding facilities must be evaluated. An environmental impact statement (EIS) should be prepared that clearly discloses the environmental consequences of, and alternatives to, the continued operation of stock holding facilities in the planning areas.

Your management plan(s) should include the following elements to address the impacts of brown-headed cowbirds:

- Remove pack stations and stables from national park lands
- Reduce stock use to the minimum amount that is necessary

Aesthetic effects—adverse impacts on visitors' experience. We are also concerned about the many aesthetic impacts that result from stock use, such as the presence of annoying bells, dust, manure, urine, and flies, and the proliferation of unsightly hoofprints, drift fences, and

⁵ See the *Water Quality Control Plan for the Central Valley Region*, the State Water Resource Control Board's Resolution No. 68-16 ("Statement of Policy with Respect to Maintaining High Quality Waters in California"), and 40 CFR § 131.12

overgrazed areas (see Absher 1979, Cole 1990, Stankey 1973, Watson et al. 1993). Most of the mitigation measures suggested above would have the added benefit of offsetting these “social” impacts. For instance, designating campsites for stock users would prevent sites used by hikers from being littered with stock manure. Tying stock and supplying feed will eliminate the need for bells and drift fences, prevent overgrazing and trampling of sensitive areas by stock, and reduce the pollution of surface waters by stock animal wastes (i.e., manure and urine).

Designation of a network of “foot travel only” trails will provide hikers with a stock-free experience (i.e., no manure or dusty trails churned by stock, etc.). Adoption of group size limits based on science (see below, especially Cole 1989 & 1990, Watson et al. 1993) will reduce the impacts of large stock groups on the experience of hikers.

Group size limits. The NPS at Yosemite has in the past taken the irresponsible, unsupportable (and illegal) position that limits on group size will only be adjusted in conjunction with surrounding land units. This ignores the mandate of the Wilderness Act and the Wild and Scenic Rivers Act to preserve wilderness and wild & scenic river values regardless of how other surrounding areas might be managed (or mismanaged). The fact that officials in the central and southern Sierra agreed on a consistent number in 1991 for maximum group sizes is no excuse to ignore the mandates of the Wilderness Act, the Wild and Scenic Rivers Act, and the Park Service’s Organic Act. This is especially true since the 15-year-old decision to allow 25 stock animals per group throughout the central/southern Sierra was adopted without following any NEPA process, and was implemented over the strong objections of hundreds of citizens and scores of conservation groups.

Further, the current group size limits have been shown to significantly and adversely affect park resources and values. In order to adequately protect Yosemite’s environment and wild & scenic river values, the group size limits must be revised downward.

Number of persons per group (on trails). Dr. David Cole, an internationally recognized research scientist, has written: **“Limits on party size must be quite low (certainly no larger than 10) to be worthwhile”** (Cole 1989). We therefore propose that group size (on trails) be limited to 10 persons, as suggested by Dr. Cole.

Number of persons per group (off trail). Large groups traveling “cross-country” cause significantly greater impacts to resources and the experience of visitors (Cole 1989 & 1990, Stankey 1973). Dr. Cole (1989) has written: **“...small parties are critical to avoid the creation of new campsites and trails in little-used places...Once a party exceeds a certain number (perhaps four to six), special care must be taken in off-trail travel.”** As suggested by Dr. Cole, group size should be limited to no more than four to six persons for all off-trail travel.

Number of stock animals per group. Dr. Cole has found that thresholds in group size that result in unacceptable impacts **“...would certainly differ between backpackers and parties with stock”** (Cole 1989). He adds that lower limits are necessary for stock parties, since they cause greater social **and** ecological impacts. Yosemite National Park must acknowledge the available research findings and conclusions, and regulate hikers and stock users according to their varying degrees of impact. The current group size regulations in effect for Yosemite’s backcountry—which employ the same limits for hikers and stock users—were arbitrarily adopted for “ease of

management.” This scheme does not comply with either the Wilderness Act, the Wild and Scenic Rivers Act, or the Park Service’s own Organic Act or wilderness management policies.

Recent research has shed light on the effects of large stock groups on the experience of wilderness users. Watson et al. (1993) documented that the average hiker in the central/southern Sierra is unacceptably affected by encountering stock groups with more than *nine* animals. Even stock users themselves are negatively affected by encounters with large groups: The average *stock user* in the central/southern Sierra is unacceptably affected by encountering groups with over *fifteen* animals (Watson et al. 1993, Table 29 & Table 10). Thus it is very clear that *twenty-five* animals in a group will degrade the character of the Tuolumne River corridor and the Tuolumne Meadows area for the majority of visitors. The Park Service must take action to prevent impairment of these areas by lowering the group size limit for stock parties.

We propose that groups be limited to no more than nine head of stock per party in the Tuolumne River corridor and Tuolumne Meadows area (*see Cole 1989 & 1990, Watson et al. 1993*), and that **all** off-trail travel by stock be prohibited.

Cross-country (off-trail) travel with stock. One very important element in Yosemite’s existing Wilderness Management Plan (WMP) is the prohibition on cross-country travel by groups with stock animals or groups over 8 persons. The plan states:

“It is Service policy to deemphasize cross-country travel by limiting such travel in Yosemite Wilderness to groups of eight people or fewer. This plan recognizes actual and potential environmental deterioration from off-trail use.”

and

“Stock must travel on designated trails or authorized stock routes and remain within one quarter mile of trails for watering, rest stops, and camping.”

This important language must be retained (and strengthened as per our comments above). We recommend against any attempt to weaken this language or to open new areas to off-trail stock use.

Two harmful loopholes in the current WMP must be addressed during the planning process for Tuolumne River/Meadows. First, the exceptions in the WMP (Appendix G) for cross-country travel by stock animals must be removed. Secondly, nowhere does the plan list or define “designated” or “established” trails. (Appendix G lists “authorized” exceptions but not the “designated” or “established” trails on which large groups are permitted). Some older maps, still in use, show trails that are no longer maintained, and which are not suitable for travel with stock or by large groups. A list or map clearly defining what trails/routes are open to travel with stock and by large groups in the Tuolumne River and Meadows areas should be addressed in these planning processes. This will make clear, to both the public and agency personnel, which routes are open and closed to travel with stock and to large groups.

We request the opportunity to review the map or list described above before it is adopted. It should be included in the draft environmental impact statements (DEISs) for these planning processes.

Summary and Conclusions

As discussed above, the three High Sierra Camps and commercial packstock enterprises are having significant, adverse impacts on the environment in both Tuolumne Meadows and within the Tuolumne River Wild & Scenic River corridor. Your plans should fully address these impacts by eliminating the HSCs, and adopting effective limits and controls on commercial packstock enterprises.

Thank you for considering the above comments, and incorporating these issues into your plans for the Tuolumne River and Tuolumne Meadows. Please contact me at the letterhead address if you have any questions about this letter. Please also send full paper copies of all environmental and decision documents for our review.

Sincerely yours,

Peter Browning
High Sierra Hikers Association

Enclosures (3): (1) *Coliform Bacteria in Sierra Nevada Wilderness Lakes and Streams: What Is the Impact of Backpackers, Pack Animals, and Cattle?* by Derlet and Carlson (2006); (2) *The Brown-headed Cowbird in the Sierra Nevada: Impacts on Native Songbirds and Possible Mitigation Measures*, by B.C. Spence; and (3) *Horses in Diapers Help Mexico's Beach Cleanup*, by Reuters, August 2003.

REFERENCES

Absher, J., and E. Absher. 1979. Sierra club wilderness outing participants and their effect on Sierra Nevada wilderness users. pp. 31-60, In: J.T. Stanley et al. (eds.) *A Report On the Wilderness Impact Study*. Sierra Club, Palo Alto, CA.

Airola, D.A. 1986. Brown-headed cowbird parasitism and habitat disturbance in the Sierra Nevada. *J Wildlife Manage* 50(4):571-575.

Ames, C.R. 1977. Wildlife conflicts in riparian management: Grazing. pp. 49-52. In: Johnson, R.R. and D.A. Jones. *Importance, Preservation and Management of Riparian Habitats*. USDA Forest Service, Gen. Tech. Rpt. RM-43. Rocky Mtn. Forest & Range Experiment Station, Ft. Collins, CO.

Armour, C. 1979. Livestock management approaches and the fisheries resource. p. 39. In: Cope, O.B. (ed). Proc. of the Forum on Grazing and Riparian/Stream Ecosystems. 94 pp. Trout Unlimited, Inc. Denver, CO.

Behnke, R.J. and R.F. Raliegh. 1978. Grazing and the riparian zone: Impact and management perspectives. pp. 184-189. In: Johnson, R.D. and J.F. McCormick. Strategies for Protection and Management of Floodplain Wetlands and other Riparian Ecosystems. 410 pp. USDA Forest Service Gen. Tech. Rpt. WO-12. Wash., D.C.

Belsky, A.J., and J.L. Gelbard. 2000. Livestock Grazing and Weed Invasions in the Arid West. A scientific report published by the Oregon Natural Desert Association. Bend, Oregon.

Bemrick, W.J. 1968. *Giardia* in North American horses. *Vet Med/SAC* 63:163-165.

Benninger, M.C. 1989. Trail corridors as conduits of movement for plant species in coniferous forests of Rocky Mountain National Park, Colorado. M.S. Thesis, Miami University.

Benninger-Truax, M.C., Vankat, J.L., and R.L. Schaefer. 1992. Trail corridors as habitat and conduits for movement of plant species in Rocky Mountain National Park, Colorado, USA. *Landscape Ecology*. 6: 269-278.

Blaser, M.J., D.N. Taylor and R.A. Feldman. 1984. Epidemiology of *Campylobacter* infections. In: Butzler, J.P. (ed.) *Campylobacter Infection in Man and Animals*. pp. 143-161. CRC Press, Inc. Boca Raton, FL.

Bohn, C.C. and J.C. Buckhouse. 1985. Some responses of riparian soils to grazing management in northeastern Oregon. *J Range Manage* 38:378-381.

Buret, A., N. denHollander, P.M. Wallis, et al. 1990. Zoonotic potential of giardiasis in domestic ruminants. *J Inf Dis* 162:231-237.

Butzler, J.P. 1984. Introduction, In: Butzler, J.P. (ed.) *Campylobacter Infection in Man and Animals*. CRC Press, Inc. Boca Raton, FL.

Campbell, J.E., and D.J. Gibson. 2001. The effect of seeds of exotic species transported via horse dung on vegetation along trail corridors. *Plant Ecology*. 157: 23-35.

Capon, A.G., J.A. Upcroft, P.F.L. Boreham, L.E. Cottis, and P.G. Bundesen. 1989. Similarities of giardia antigens derived from human and animal sources. *International Journal for Parasitology* 19(1):91-98.

Cole, D.N. 1977. Man's impact on wilderness vegetation: an example from the Eagle Cap Wilderness, northeastern Oregon. Ph.D. diss. Univ. Oregon, Eugene. 307 p.

Cole, D.N. 1989. Low-impact recreational practices for wilderness and backcountry. USDA Forest Service, Intermountain Research Station, Gen. Tech. Rpt. INT-265. Ogden, UT.

Cole, D.N. 1990. Ecological impacts of wilderness recreation and their management. pp. 425-462, In: Hendee, J.C., et al. (eds.) *Wilderness Management*. Second edition. North American Press, Golden, CO.

Dale, D., and T. Weaver. 1974. Trampling effects on vegetation of the trail corridors of north Rocky Mountain forests. *J Appl Ecol* 11:767-772.

Davies, R.B. and C.P. Hibler. 1979. Animal reservoirs and cross-species transmission of *Giardia*. pp. 104-126. In: *Waterborne Transmission of Giardiasis*. U.S. Environmental Protection Agency (EPA-600/9-79-001), Cincinnati, OH.

Davis, G.A. 1977. Management alternatives for the riparian habitat in the Southwest. pp. 59-67. In: Johnson, R.R. and D.A. Jones. *Importance, Preservation and Management of Riparian Habitats*, USDA Forest Service Gen. Tech. Rpt. RM-43. 217 pp. Rocky Mtn. Forest & Range Experiment Sta., Fort Collins, CO.

Derlet, R.W., and J.R. Carlson. 2002. An Analysis of Human Pathogens Found in Horse/Mule Manure Along the John Muir Trail in Kings Canyon and Sequoia and Yosemite National Parks. *Wilderness and Environmental Medicine* 13:113-118.

Derlet, R.W., and J.R. Carlson. 2006. Coliform Bacteria in Sierra Nevada Wilderness Lakes and Streams: What Is the Impact of Backpackers, Pack Animals, and Cattle? *Wilderness and Environmental Medicine* 17:15-20.

Duff, D.A. 1979. Riparian habitat recovery on Big Creek, Rich County, UT. p. 91. In: Cope, O.B. (ed). Proc. of the Forum on Grazing and Riparian/Stream Ecosystems. 94 pp. Trout Unlimited, Inc. Denver, CO.

Faubert, G.M. 1988. Evidence that giardiasis is a zoonosis. *Parasitology Today* 4(3):66-68.

Fenner, M. 1985. Seed ecology. Chapman & Hall, London.

Fleischer, R.C., A.P. Smith and S.I. Rothstein. 1987. Temporal and age-related variation in the laying rate of the parasitic brown-headed cowbird in the eastern Sierra Nevada, California. *Can J Zool* 65:2724-2730.

Frissell, S.S. 1973. The impact of wilderness visitors on natural ecosystems. Unpubl. rep., USDA Forest Service, Forest Science Laboratory, Missoula, MT. 60 p.

Graber, D.M. 1996. Status of Terrestrial Vertebrates. pp. 709-734, In: Sierra Nevada Ecosystem Project: Final report to Congress, vol. II, Assessments and scientific basis for management options. University of California, Davis, Centers for Water and Wildland Resources, 1996.

Hammit, W.E., and D.N. Cole. 1987. Wildland Recreation: Ecology and management. John Wiley & Sons, New York.

Harmon, G.W., and F.D. Kiem. 1934. The percentage and viability of weed seeds recovered in the feces of farm animals and their longevity when buried in manure. *J. Am. Soc. Agron.* 26: 762-767.

Harper, J.L. 1977. Population biology of plants. Academic Press, London.

Heady, H.F. 1954. Viable seed recovered from fecal pellets of sheep and deer. *Journal of Range Management*. 7: 259-261.

Hendee, J.C., G.H. Stankey, and R.C. Lucas. 1990. *Wilderness Management*. Second edition. International Wilderness Leadership Foundation in cooperation with the USDA Forest Service. North American Press, Golden, CO.

Isaac-Renton, J.L., C. Cordeiro, C. Sarafis, and H. Shahriari. 1993. Characterization of giardia duodenalis isolates from a waterborne outbreak. *Journal of Infectious Diseases* 167:431-440.

Janzen, D.H. 1981. *Enterlobium cyclocarpun* seed passage rate and survival in horses, Costa Rican pleistocene seed dispersal agents. *Ecology*. 62: 593-601.

Janzen, D.H. 1982. Differential seed survival and passage rates in cows and horses. *Oikos*. 38: 150-156.

Kasprzak, W., and Z. Pawlowski. 1989. Zoonotic aspects of giardiasis: a review. *Vet Parasitol* 32:101-108.

Kauffman, J.B., W.C. Krueger, and M. Vaura. 1983. Impacts of cattle on streambanks in northeastern Oregon. *J Range Manage* 36:683-685.

Kauffman, J.B., and W.C. Krueger. 1984. Livestock impacts on riparian ecosystems and streamside management implications. *J Range Manage* 37:430-438.

Kirkpatrick, C.E. 1989. Giardiasis in large animals. *Compend Contin Educ Pract Vet* 11:80-84.

Kirkpatrick, C.E., and D.L. Skand. 1985. Giardiasis in a horse. *JAVMA* 187:163-164.

Knudson, T. 1991. Roads, cattle damage Sierra: Logging, grazing cause soil erosion, watershed problems. *Oakland Tribune*. 15 December 1991.

Kuss, F.R., A.R. Graefe, and L. Loomis. 1986. Plant and soil responses to wilderness recreation: a synthesis of previous research. pp. 129-137, In: Lucas, R.C. (compiler) *Proceedings—National Wilderness Research Conference: Current Research*. USDA Forest Service, Intermountain Research Station, Ogden UT. General Technical Report INT-212.

Laing, C.C. 1961. A report on the effect of visitors on the natural landscape in the vicinity of Lake Solitude, Grand Teton National Park, Wyoming. Unpubl. rep. USDI National Park Service, Grand Teton National Park. 62 p.

Landsberg, J., B. Logan, and D. Shorthouse. 2001. Horse riding in urban conservation areas: Reviewing scientific evidence to guide management. *Ecological Management & Restoration*. 2(1): 36-46.

LeChevallier, M.W., W.D. Norton, and R.G. Lee. 1991. *Appl Environ Microbiol* 57:2610-2616.

Manahan, F.F. 1970. Diarrhoea in horses with particular reference to a chronic diarrhoea syndrome. *Aust Vet J* 46:231-234.

Manser, P.A., and R.W. Dalziel. 1985. A survey of *Campylobacter* in animals. *J Hyg (Camb)* 95:15-21.

Marlow, C.B., and T.M. Pogacnik. 1985. Time of grazing and cattle-induced damage to streambanks. pp. 279-284. In: Johnson, R.R., Riparian Ecosystems and their Management: Reconciling Conflicting Uses. First North American Riparian Conf. USDA Forest Service. Gen. Tech. Rpt. RM-120. 523 pp. Rocky Mtn. Forest & Range Exp. Sta., Fort Collins, CO.

McCully, W.G. 1951. Recovery and viability of Macartney rose seeds fed to cattle. *Journal of Range Management*. 4: 101-106.

McQuaid-Cook, J. 1978. Effects of hikers and horses on mountain trails. *J Envir Mgm* 6:209-212.

Meecham, W.R., and W.S. Platts. 1978. Livestock grazing and the aquatic environment. *J Soil & Water Conser* 33(6):274-278.

Merkle, J. 1963. Ecological studies of the Amphitheater and Surprise Lakes cirque in the Teton Mountains, Wyoming. Unpubl. rep., National Park Service, Grand Teton National Park, 25 p.

Meyer, E.A. 1988. Waterborne *Giardia* and *Cryptosporidium*. *Parasitology Today* 4(7):200-201.

Miller, T.C. East Bay water quality threat? *The Daily Californian*. 16 December 1991.

Mosconi, S.L., and R.L. Hutto. 1982. The effect of grazing on the land birds of a western Montana riparian habitat. pp. 221-233. In: Peek, J.M., and P.D. Dalke (eds). Wildlife-Livestock Relationships Symposium. Proc. 10. 614 pp. Univ. of Idaho. Forest, Wildlife and Range Exp. Sta., Moscow, ID.

Nagy, J.A.S., and G.W. Scotter. 1974. A quantitative assessment of the effects of human and horse trampling on natural areas. Waterton Lake National Park. Unpubl. rep., Canadian Wildlife Service, Edmonton, Alberta. 145 p.

Neuman, M.J. 1990. Past and present conditions of backcountry meadows in Sequoia and Kings Canyon National Parks. Second edition. Unpubl. rep., USDI National Park Service, Sequoia & Kings Canyon NP's, Three Rivers, CA. 21 December 1990.

Neuman, M.J. 1991a. Accomplishments of the stock use and meadow monitoring program in 1990. Unpubl. rep., USDI National Park Service, Sequoia & Kings Canyon NP's, Three Rivers, CA. 18 January 1991.

Neuman, M.J. 1991b. 1991 Accomplishments of the stock use and meadow monitoring program. Unpubl. rep., USDI National Park Service, Sequoia & Kings Canyon NP's, Three Rivers, CA. 1 November 1991.

Piggin, C.M. 1978. Dispersal of *Echium plantagineum* L. by sheep. *Weed Res.* 18: 155-160.

Platts, W.S. 1979. Livestock grazing and riparian/stream ecosystems. pp. 39-45. In: Cope, O.B. (ed). Proc. of the Forum on Grazing and Riparian/Stream Ecosystems. 94 pp. Trout Unlimited, Inc. Denver, CO.

Platts, W.S. 1981. Effects of sheep grazing on a riparian-stream environment. USDA Forest Service Research Note INT-307. Intermountain Forest & Range Exp. Sta., Ogden, UT.

Platts, W.S., and F.J. Wagstaff. 1984. Fencing to control livestock grazing on riparian habitats along streams: Is it a viable alternative? *NAJMDP* 4:266-272.

Quinn, L., M. Kolipinski, and S. Ghosh. 2006. Horses as vectors for weed invasion in California's National Parks: The promise of weed-free feed. Presented at: 59th Annual Meeting of the Western Society of Weed Science, March 14-16, 2006. Reno, NV.

Ridley, H.N. 1930. Dispersal of Plants Throughout the World. Reeve, Ashford, UK.

Rosquist, A. 1984. *Giardia* source investigation in Rattlesnake Creek, July 5 - October 31, 1983. Unpubl. memo. Lolo National Forest, Missoula, MT.

Rothstein, S.I., J. Verner, and E. Stevens. 1980. Range expansion and diurnal changes in dispersion of the brown-headed cowbird in the Sierra Nevada. *Auk* 97:253-267.

Rush, B.A., P.A. Chapman, and R.W. Ineson. 1987. *Lancet* 2:632-633.

Saeed, A.M., N.V. Harris, and R.F. DiGiacomo. 1993. The role of exposure to animals in the etiology of campylobacter jejuni/coli enteritis. *American J of Epidemiology* 137(1):108-114.

Siekert, R.E., Q.D. Skinner, M.A. Smith, J.L. Doad, and J.D. Rodgers. 1985. Channel response of an ephemeral stream in Wyoming to selected grazing treatments. pp. 276-278. In: Johnson, R.R. Riparian Ecosystems and their Management: Reconciling Conflicting Uses. First North American Riparian Conf. USDA Forest Service Gen. Tech. Rpt. RM-120. 523 pp. Rocky Mtn. Forest & Range Exp. Sta., Fort Collins, CO.

St. Jean, G., Y. Couture, P. Dubreuil, and J.L. Frechette. 1987. Diagnosis of *Giardia* infection in 14 calves. *JAVMA* 191(7):831-832.

St John-Sweeting, R.S., and K.A. Morris. 1991. Seed transmission through the digestive tract of the horse. In: *Plant Invasions—The Incidence of Environmental Weeds in Australia*. pp. 170–172. Kowari 2, Australian National Parks and Wildlife Service, Canberra.

Stankey, G.H. 1973. Visitor perception of wilderness recreation carrying capacity. Res. Pap. INT-142. USDA Forest Service, Intermountain Forest And Range Experiment Station, Ogden UT. 61 p.

Stanley, J.T., H.T. Harvey, and R.J. Hartesveldt (eds.). 1979. *A Report On the Wilderness Impact Study*. Sierra Club, Palo Alto, CA. pp. 17, 201-202.

Strand, S. 1972. An investigation of the relationship of pack stock to some aspects of meadow ecology for seven meadows in Kings Canyon National Park. M.A. thesis. Calif. State Univ., San Jose. 125 p.

Strand, S. 1979a. The impact of stock on wilderness meadows in Sequoia and Kings Canyon National Parks. pp. 77-87. In: Stanley, J.T. et al. (eds.), *A Report On the Wilderness Impact Study*. Sierra Club, Palo Alto, CA.

Strand, S. 1979b. Recovery of Sierran meadows after trampling by pack stock. pp. 88-93. In: Stanley, J.T. et al. (eds.), *A Report On the Wilderness Impact Study*. Sierra Club, Palo Alto, CA.

Strand, S. 1979c. Pack stock management in the high Sierra. pp. 209. In: Stanley, J.T. et al. (eds.), *A Report On the Wilderness Impact Study*. Sierra Club, Palo Alto, CA.

Stranden, A.M., J. Eckert, and P. Kohler. 1990. Electrophoretic characterization of giardia isolated from humans, cattle, sheep, and a dog in Switzerland. *J Parasitol* 76(5):660-668.

Suk, T.J. 1983. Investigation of animal hosts for *Giardia* spp. in California's Sierra Nevada mountains. USDI National Park Serv., Coop. Park Studies Unit, Institute of Ecology, Univ. CA at Davis. Tech. Rpt. No. 11.

Suk, T.J., J.L. Riggs, and B.C. Nelson. 1986. Water contamination with *Giardia* in backcountry areas. In: Proc. of the National Wilderness Research Conf. Gen. Tech. Rpt. INT-212. USDA Forest Service Intermountain Research Sta., Ogden, UT.

Sumner, L., and R.M. Leonard. 1947. Protecting mountain meadows. *Sierra Club Bulletin* 32(5):53-69.

Taylor, D.N., K.T. McDermott, J.R. Little, J.G. Wells, M.J. Blaser. 1983. *Campylobacter* enteritis from untreated water in the Rocky Mountains. *Ann Intern Med* 99:38-40.

Taylor, D.M. 1986. Effects of cattle grazing on passerine birds nesting in riparian habitat. *J Range Manage* 39:254-258.

Thomas, J.W., C. Maser, and J.E. Rodnick. 1979. Riparian zones in managed rangelands: their importance to wildlife. pp. 21-30. In: Cope, O.B. (ed). Proc. of the Forum on Grazing and Riparian/Stream Ecosystems. 94 pp. Trout Unlimited, Inc., Denver, CO.

Trottier, G.C., and G.W. Scotter. 1975. Backcountry management studies, the Egypt Block, Banff National Park. Unpubl. rep., Canadian Wildlife Service, Edmonton, Alberta. 178 p.

Upcroft, J.A., and P. Upcroft. 1994. Two distinct varieties of giardia in a mixed infection from a single patient. *J Euk Microbiol* 41(3):189-194.

Verner, J. and L.V. Ritter. 1983. Current status of the brown-headed cowbird in the Sierra National Forest. *Auk* 100:355-368.

Watson, A., M.J. Niccolucci, and D.R. Williams. 1993. Hikers and recreational stock users: predicting and managing recreation conflicts in three wildernesses. Research Paper INT-468. USDA Forest Service, Intermountain Research Station. Ogden, UT.

Weaver, V., and R. Adams. 1996. Horses as vectors in the dispersal of weeds into native vegetation. Proceedings of the Eleventh Australian Weeds Conference. pp. 383–397.

Weaver, T., and D. Dale. 1978. Trampling effects of hikers, motorcycles and horses in meadows and forests. *J Appl Ecol* 15:451-457.

Weaver, T., D. Dale, and E. Hartley. 1979. The relationship of trail condition to use, vegetation, user, slope, season and time. pp. 94-100, In: Ittner, R., et al. (eds.), *Recreational Impact on Wildlands* conference proceedings, Oct. 27-29, 1978, USDA Forest Service, Pacific Northwest Region, R-6-001-1979.

Welch, D. 1985. Studies in the grazing of heather moorland in northeast Scotland; IV. Seed dispersal and plant establishment in dung. *Journal of Applied Ecology*. 22: 461-472.

Weniger, B.G., et al. *Am J Public Health* 73:868-872.

Whitson, P.D. 1974. The impact of human use upon the Chisos Basin and adjacent lands. USDI National Park Service Science Monograph Series 4. Gov. Print. Office, Washington, D.C. 92 p.

Whittaker, P.L. 1978. Comparison of surface impact by hiking and horseback riding in the Great Smoky Mountains National Park. Manage. Rep. 24. USDI National Park Service, Southeast Region. 32 p.

Wilson, J.P., and J.P Seney. 1994. Erosional impact of hikers, horses, motorcycles, and off-road bicycles on mountain trails in Montana. *Mountain Research and Development* 14(1):77-88.

Winegar, H.H. 1977. Camp Creek channel fencing: plant, wildlife, and soil and water response. *Rangeman's Journal* 4:10-12.

Xiao, L., R.P. Herd, and D.M. Rings. 1993. *Vet Parasitol* 51:41-48.